

## **Marine Mammals: Hearing and Echolocation at Coconut Island**

Paul E. Nachtigall  
Marine Mammal Research Program  
P.O. Box 1106  
Kailua HI 96734

phone: (808) 247-5297    fax: (808) 247-5831    email: [nachtiga@hawaii.edu](mailto:nachtiga@hawaii.edu)

Award Number: N000140811160  
<http://www.hawaii.edu/HIMB/>

### **LONG-TERM GOALS**

1. Broaden the baseline of hearing measures of marine mammals by increasing the number of animals and species measured. The effects of sound on wild populations of animals can best be determined if baseline hearing measures are known.
2. Develop an understanding of the basic processes of odontocete echolocation for ecosystem management. Echolocation is the principal sense for odontocete foraging and its understanding is therefore crucial for estimating the effects of sound on foraging and populations and ecosystem management.
3. Examine the control of hearing during echolocation.
4. Improve the measurement of marine mammal hearing by developing and refining hearing procedures particularly those that will rapidly measure the hearing of stranded and temporarily caught animals.
5. Comparatively examine the basic hearing mechanisms of marine mammal species.
6. Comparatively examine the hearing during echolocation of species in addition to that of the false killer whale.
7. Examine automatic gain control mechanisms in odontocete echolocation.

### **OBJECTIVES**

Marine Mammal sensory systems have evolved to effectively use acoustic energy in the oceans. Our objectives are to develop a basic understanding of hearing and echolocation so that knowledge can then be applied to the solution of practical problems as they arise. The most basic hearing measurement is the audiogram which is a series of thresholds across frequencies. It basically describes the hearing of an organism. Audiograms are the most basic of the hearing measures and are essential for describing the audiometrics of a species of animals. Of the 85 species of dolphins and whales we

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>2012</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Marine Mammals: Hearing and Echolocation at Coconut Island</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Marine Mammal Research Program P.O. Box 1106 Kailua, HI 96734</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>SAR</b>	18. NUMBER OF PAGES <b>8</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

now have audiograms on 17 species. Audiograms on additional cetacean, and other marine mammal, species may be obtained from stranded animals, from animals in captive display situations, and from catch and release scenarios. We intend to obtain as many valid audiograms as possible as we seek new opportunities in new situations. Most marine mammal audiograms are obtained on individuals and published individually. Population estimates obtain increased validity with increased numbers of measurements. Other hearing measures such as directionality of hearing, and the mechanisms underlying that directionality, are also very important and little is known on most marine mammals. These measures will also be obtained whenever possible.

Most of our initial audiometric work measured hearing using behavioral responses (Nachtigall et al, 2000). Measures of auditory evoked potentials (AEP) produce the benefit of being obtained rapidly without requiring captivity or lengthy training. Our work (Yuen et al, 2005) shows that the two procedures, while not producing exactly the same results, are certainly comparable. So, we intend to continue using AEP measures to measure the hearing of new species and to continue to measure hearing in other situations.

Our initial work on temporary threshold shifts (Nachtigall et al 2003, 2004) with exposures up to 50 minutes combined with shorter term exposures led to an equal energy hypothesis in which it was assumed that the amount of TTS was dependent on the amount of energy received relatively independent of the time of exposure. An objective of our recent work (Mooney et al, 2009a) has been to examine whether the equal energy hypothesis is valid for short exposure times and to examine the direct effects of the Navy 53C on the hearing of the bottlenose dolphin (Mooney et al, 2009b).

While much was known about outgoing dolphin echolocation signals, little was known about what animals heard while they echolocated until we developed a procedure to measure AEPs during active echolocation experiments (Supin et al. 2003 ). Our current objective is to examine the automatic gain control of hearing during echolocation with targets present and absent (Supin et al 2011) and then to further explore whether or not the dolphins and whales can change their hearing in response to loud sounds being presented. Our work with echolocation demonstrated that hearing sensitivity changed with echolocation task and situation. Can hearing sensitivity also change if an arbitrary sound is presented as a warning prior to loud sound? We are interested in basic echolocation processes and the interaction of hearing an echolocation. What specifically contributes to the outstanding echolocation discrimination capability of the odontocetes. How important is high frequency hearing to echolocation discrimination.

The ability to obtain hearing data on new species requires opportunistic motivated action. A permit from the NMFS to test the hearing of stranded animals, and keep in touch with the stranding networks in the US, allows us opportunities for testing stranded animals when they become available. Animals in public display facilities also occasionally become available for hearing examination especially if the audiometric tests are conducted for short periods of time like those for AEP measures. The AEP measures can also be used on boats so that temporarily caught animals can be tested. All of these approaches allow us to increase the species and the number of animals tested. We work closely with Alexander Ya. Supin from the Russian Academy of Sciences to test the hearing of new species of animals, especially those that require a new technique or adaptations to new procedures.

We primarily use the envelope following response (EFR) auditory evoked potential approach in which we present amplitude modulated sounds to the animals and monitor the brain wave patterns in response to the amplitude modulation rate.

Hearing is examined during echolocation by measuring the hearing when the outgoing pulse is produced and when the echo is returned by measuring the brain response timed from the outgoing pulse. Target strength is varied by changing the distance and target strength of the targets. The animal is required to echolocate and report the presence or absence of the target and AEP recordings of its hearing while doing that are the important dependent variable. Our work on the False killer whale shows precise control of hearing during echolocation.

## **APPROACH**

The ability to obtain hearing data on new species requires opportunistic motivated action. A permit from the NMFS to test the hearing of stranded animals, and keep in touch with the stranding networks in the US, allows us opportunities for testing stranded animals when they become available. Animals in public display facilities also occasionally become available for hearing examination especially if the audiometric tests are conducted for short periods of time like those for AEP measures. The AEP measures can also be used on boats so that temporarily caught animals can be tested. All of these approaches allow us to increase the species and the number of animals tested. We work closely with Alexander Ya. Supin from the Russian Academy of Sciences to test the hearing of new species of animals, especially those that require a new technique or adaptations to new procedures.

We primarily use the envelope following response (EFR) auditory evoked potential approach in which we present amplitude modulated sounds to the animals and monitor the brain wave patterns in response to the amplitude modulation rate. We also use the AEP approach of simply examining the brain wave patterns to impulsive sounds monitored with time.

Hearing is examined during echolocation by measuring the hearing when the outgoing pulse is produced and when the echo is returned by measuring the brain response timed from the outgoing pulse. Target strength is varied by changing the distance and target strength of the targets. The animal is required to echolocate and report the presence or absence of the target and AEP recordings of its hearing while doing that are the important dependent variable. Our work on the False killer whale shows precise control of hearing during echolocation. Recent work (Li et al, 2011) shows the same processes occur with other odontocetes like the bottlenose dolphin. More recent work (Linnenschmidt 2011, 2012) shows similar processes ongoing with the harbor porpoise (*Phocoena phocoena*).

Hearing is a remarkably evolved process in the odontocetes. Efforts in progress show that hearing directionality is dependent on frequency. Hearing is measured with evoked potentials as different frequency sounds are presented from a variety of directions.

Outgoing echolocation clicks are also measured as a function of task. Although many people have assumed that the melon functions as a ‘focusing’ device, little published empirical data support the notion. Measures of echolocating animals in different situations demonstrate empirical changes in the outgoing beam pattern likely caused by focusing of the melon.

## **WORK COMPLETED**

Completed and published the audiogram of a stranded beaked whale *Mesoplodon densirostris* and showed that its best hearing was the same as its published echolocation return frequencies at 40 to 50 kHz.

Continued the measurement of hearing during echolocation on the false killer whale demonstrating the ability of the whale to change hearing during echolocation. Demonstrated that there was an expectation component in changing the echolocation pulses (Supin et al, 2012).

Completed comparative measurements of hearing during echolocation on the bottlenose dolphin and the harbor porpoise. Both demonstrate control of hearing in a manner similar to the false killer whale (Li et al, 2011 & 2012, Linnenschmidt et al, 2011 & 2012).

Tested the comparative hearing pathways of the bottlenose dolphin and the false killer whale.

Examined the outgoing beam focusing during echolocation of the false killer whale and the harbor porpoise. (Kloepper et al, 2011 & 2012).

## **RESULTS**

The formerly stranded beaked whale heard relatively well in the midfrequency sonar range (5.6 kHz) at 79 dB.

Published results indicating that a formerly stranded long finned pilot whale and a recently stranded pygmy killer whale showed audiograms with high frequency hearing (Pacini et al, 2011 & 2012).

Found that there is an automatic gain control in the hearing system of an echolocating false killer whale. There are at least three mechanisms of automatic gain control in odontocete echolocation – echolocation and hearing are a very dynamic process. Began experiments to determine whether the control of hearing was confined to echolocation.

Found that it is unlikely that the control of hearing during echolocation is accomplished via a stapedial mechanism in the same way that it is controlled by bats.

Found that hearing pathways differ between the false killer whale, the bottlenose dolphin and the beluga whale. Initial work indicates that the hearing pathway of sound in the dolphin may not be the traditionally assumed lower jaw acoustic window route. Additional work and modeling is necessary.

## **IMPACT/APPLICATIONS**

High frequency sonar issues may be an important future area . Exposure to loud sound effects echolocation. Long term exposure can result in decrements in echolocation and foraging. Long-term ecosystem effects are likely.

We still do not know the hearing range and sensitivity of the mysticete whales. We presume they hear low frequencies, but perhaps like the polar bears, they hear much higher frequencies than we assume. If that is true, it is reasonable to be concerned about the effects of sound on mysticete whales. The mysticete whale hearing issue remains and important one to be resolved. We continue our efforts to obtain a subject for hearing examination and have written the mysticete whales into our new permit.

The false killer whale's, the bottlenose dolphin's and the harbour porpoise's overall hearing changes while it echolocates. Does it control its hearing in a voluntary way? Can this be used for mitigation? Can a warning sound be given that allows the animal to change its hearing?

All odontocetes are not alike. The audiogram of the beaked whale differs greatly from other odontocetes. We assume that the Sperm whale will differ even more. The idea of using the bottlenose dolphin as a template for all odontocete hearing is not a reasonable assumption for modeling or for regulatory purposes.

Belugas and false killer whales hear directionally very differently than the bottlenose dolphin. We must be cautious when we extrapolate from species to species. More work is needed. A combination of modeling and empirical hearing data is necessary.

## TRANSITIONS

The audiogram on the beaked whale (*Mesoplodon densirostris*) provides the Fleet with an actual good measure of the hearing thresholds of beaked whales in the mid-frequency sonar frequency. This data is essential for planning fleet sonar exercises and for determining the number of ‘takes’ in permits where beaked whales are likely to occur.

Our results on Hearing and also TTS are used by the U.S. Naval Fleets in their requests for LOAs and permits for Training. Our data are also cited by NOAA, NMFS in their Biological opinions required under the Endangered Species Act for issuance of opinions regarding Fleet Training and in Navy and NMFS NEPA requirements. The SPAWARSCEN in San Diego California uses the basic procedure that we developed to test dolphin hearing with auditory evoked potentials in research and also to measure the hearing of Navy systems animals.

## RELATED PROJECTS

In a related project, funded via the University of Southern Denmark we are working with a new post-doctoral fellow, Meike Linnenschmidt and her professor Magnus Wahlberg, to expand the effort of examination of hearing during echolocation looking at harbour porpoises. The results indicate that the harbor porpoise hears during echolocation in a manner similar to the false killer whale. Two manuscript has been written and published by Ms. Linnenschmidt. Our (former)graduate student, Dr. Laura Kloepper received a Jnl Exp Biol fellowship to travel to Denmark to work with Drs. Wahlberg and Linnenschmidt. Newly published work (Linnenschmidt et al 2012) demonstrated control of hearing and outgoing echolocation pulse changes during echolocation.

We continue to examine the directionality of hearing and work with the modeling efforts of both Dr. Ted Cranford at San Diego State University and Dr. Darlene Ketten and (former student) Dr. Aran Mooney at Woods Hole Oceanographic Institution. One of our graduate students, Kristen Taylor is specifically examining the head related transfer function of the bottlenose dolphin in collaboration with Dr. Cranford.

## REFERENCES

- Kloepper, L. N., Nachtigall, P. E., Donahue, M. J., and Breese,(2012) M. Active echolocation beam focusing in the false killer whale (*Pseudorca crassidens*) *Jnl Exp Biol* 1306-1312.
- Kloepper, L.N., Nachtigall, P.E., Quintos, C., & Vlachos, S. (2012) Single-lobed frequency-dependent Beam shape in an echolocating false killer whale (*Pseudorca crassidens*) *J Acoust Soc Am* (in press).

- Li, S. Nachtigall, P.E., and Marlee Breese. (2011) Dolphin hearing during echolocation: evoked potential responses in an Atlantic bottlenose dolphin (*Tursiops truncatus*). *Jnl Exp Biol* 214, 2027-2035 .
- Li S, Nachtigall PE, Breese M, Supin AY (2012) Hearing Sensation Levels of Emitted Biosonar Clicks in an Echolocating Atlantic Bottlenose Dolphin. *PLoS ONE* 7(1): e29793. doi:10.1371/journal.pone.0029793
- Linnenschmidt M., Kloepper, L. N. , M. Wahlberg, M. and Nachtigall, P.E. (2012) Extremely rapid gain control regulation in the biosonar of a harbour porpoise. *Naturwissenschaften* DOI 10.1007/s00114-012-0948-7. *Published online*
- Linneschmidt, M., Beedholm, K., Wahlberg, M, Hojer-Kristensen, J and Nachtigall, P.E. Keeping returns optimal: Gain control elicited by dynamic hearing thresholds in a harbour porpoise. (2012) *Biol Letters Proceedings of the Royal Society B*. doi:10.1098/rspd.2011.2465. *published online*
- Mooney T.A., Nachtigall, P.E. Breese, M. Vlachos, S. and Au, W.L. (2009a), Predicting temporary threshold shifts in a bottlenose dolphin (*Tursiops truncatus*): the effects of noise level and duration. *J Acous Soc Am* , 125, 1816-1826. (published, refereed)
- Mooney T.A., Nachtigall, P.E. and Vlachos, S. (2009b) Sonar induced temporary hearing loss in dolphins. *Biology Letters* , 5, 565-567 (published, refereed)
- Nachtigall, P.E., Lemonds, D.W., and Roitblat, H. L. (2000) Psychoacoustic Studies of Whale and Dolphin Hearing. In: Au, W.W.L, Popper, A.N. and Fay R.J. (eds) *Hearing By Whales*, Springer-Verlag, New York pp. 330-364.
- Nachtigall, P.E., Pawlowski, J. L., and Au W.W.L. (2003) Temporary threshold shifts and recovery following noise exposure in the Atlantic Bottlenosed dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America* 113(6) 3425-3429
- Nachtigall, P.E., Supin, A.Ya. Pawloski, J. L and Au, W.W.L. (2004) Temporary threshold shifts after noise exposure in a bottlenosed dolphin (*Tursiops truncatus*) measured using evoked auditory potentials. *Marine Mammal Science*, 20(4), 673-687
- Pacini, A.F., Nachtigall, P.E., Quintos, C., Schofield, D. Look, D.A., Levine, G. and Turner, J. (2011) Audiogram of a stranded Blainville's beaked whale (*Mesoplodon densirostris*) measured using auditory evoked potentials. *Jnl Exp Biol* 214, 2409-2415.
- Pacini, A., Nachtigall, P.E. and Kloepper, L.K. (2012) Portable auditory evoked potential system to assess odontocete hearing. Popper, A. N. and Hawkins, A. eds. *Effects of Noise on Aquatic Life*. Springer Science+Business Media, LLC, New York. 225-227.
- Supin, A.Ya., Nachtigall, P.E., Pawloski, J.L., and Au, W.W.L. (2003) Evoked potential recording during echolocation in a false killer whale (*Pseudorca crassidens*). *Journal of the Acoustical Society of America*, 113(5), 2408-2411

- Supin, A. Ya., Nachtigall, P.E., and Breese, M. (2009) Forward masking based gain control in odontocete biosonar: an evoked-potential study. *J Acous Soc Am*
- Yuen, M.E., Nachtigall, P.E., and Supin, A.Ya., and Breese, M. (2005) Behavioral and auditory evoked potential audiograms of a false killer whale (*Pseudorca crassidens*). *Journal of the Acoustical Society of America*, 118 (4) 2688-2695
- Supin, A. Ya., Nachtigall, P.E. & Breese, M. (2012) A whale better adjusts the biosonar to ordered rather than to random changes in the echo parameters. *J Acoust Soc Am*

## **PUBLICATIONS -Refereed**

### **In Press:**

- Supin, A. Ya., Nachtigall, P.E. & Breese, M. (2012) A whale better adjusts the biosonar to ordered rather than to random changes in the echo parameters. *J Acoust Soc Am*
- Kloepper, L.N., Nachtigall, P.E., Quintos, C., & Vlachos, S. (2012) Single-lobed frequency-dependent Beam shape in an echolocating false killer whale (*Pseudorca crassidens*) *J Acoust Soc Am*

### **Published:**

- Lemonds, D.W., Au, W.W.L., Vlachos, S.A., and Nachtigall, P.E. (2012) High Frequency auditory filter shape for the Atlantic bottlenose dolphin. *Jnl Acous Soc Am*, 132, 1222-1228
- Linnenschmidt M., Kloepper, L. N. , M. Wahlberg, M. and Nachtigall, P.E. (2012) Extremely rapid gain control regulation in the biosonar of a harbour porpoise. *Naturwissenschaften* DOI 10.1007/s00114-012-0948-7. *Published online*
- Kloepper, L. N., Nachtigall, P. E., Donahue, M. J., and Breese, (2012) M. Active echolocation beam focusing in the false killer whale (*Pseudorca crassidens*) *Jnl Exp Biol* 1306-1312
- Linneschmidt, M., Beedholm, K., Wahlberg, M, Hojer-Kristensen, J and Nachtigall, P.E. Keeping returns optimal: Gain control elicited by dynamic hearing thresholds in a harbour porpoise. (2012) *Biol Letters. Proceedings of the Royal Society B.* doi:10.1098/rspd.2011.2465. *published online*
- Li S, Nachtigall PE, Breese M, Supin AY (2012) Hearing Sensation Levels of Emitted Biosonar Clicks in an Echolocating Atlantic Bottlenose Dolphin. *PLoS ONE* 7(1): e29793. doi:10.1371/journal.pone.0029793
- Mooney, T.A., Hanlon, R. Madsen, P.T., Christensen-Dalsgaard, J. Ketten, D.R & Nachtigall, P.E. (2012), Potential for Sound Sensitivity in Cephalopods. Popper, A. N. and Hawkins, A. eds. *Effects of Noise on Aquatic Life*. Springer Science+Business Media, LLC, New York. 125-128.
- Pacini, A., Nachtigall, P.E. and Kloepper, L.K. (2012) Portable auditory evoked potential system to assess odontocete hearing. Popper, A. N. and Hawkins, A. eds. *Effects of Noise on Aquatic Life*. Springer Science+Business Media, LLC, New York. 225-227.

- Nachtigall, P.E., Supin, A.Ya., and Breese, M. Hearing, Noise and Echolocating Odontocetes. Popper, A. N. and Hawkins, A. eds. (2012). *Effects of Noise on Aquatic Life*. Springer Science+Business Media, LLC, New York. 49-51.
- Li, S., Nachtigall, P.E., and Breese, M. Auditory brainstem responses associated with echolocation in an Atlantic bottlenose dolphin (*Tursiops truncatus*). Popper, A. N. and Hawkins, A. eds. (2012). *Effects of Noise on Aquatic Life*. Springer Science+Business Media, LLC, New York. 45-47.
- Ibsen, S., Krause-Nehring, J., Nachtigall, P.E., Au, W.W.L, and Breese, M. (2011) Similarities in echolocation strategy and click characteristics between a *Pseudorca crassidens* and a *Tursiops truncatus*, *J Acoust Soc Am* 130, 3085-3089.
- Lemons, D.W., Kloepper, L.N., Nachtigall, P.E., Au, W.W.L, Vlachos, S.A., and Branstetter, B.K. (2011) A re-evaluation of auditory filter shape in delphinids odontocetes: evidence of constant-bandwidth filters. *J Acous Soc Am* 130, 3107-3114
- Supin, A.Ya., Nachtigall, P.E. and Breese, M. (2011) Interaction of emitted sonar pulses and simulated echoes in a false killer whale: An evoked-potential study. *J Acous Soc Am*, 130, 1711-1720
- Pacini, A.F., Nachtigall, P.E., Quintos, C., Schofield, D. Look, D.A., Levine, G. and Turner, J. (2011) Audiogram of a stranded Blainville's beaked whale (*Mesoplodon densirostris*) measured using auditory evoked potentials. *Jnl Exp Biol* 214, 2409-2415.
- Li, S. Nachtigall, P.E., and Marlee Breese. (2011) Dolphin hearing during echolocation: evoked potential responses in an Atlantic bottlenose dolphin (*Tursiops truncatus*). *Jnl Exp Biol* 214, 2027-2035

## **HONORS/AWARDS/PRIZES**

- Graduate student Laura Kloepper won first prize for the best student paper for her talk in Animal BioAcoustics at the Joint Meeting of the Acoustical Societies of China and America in Hong Kong And second prize overall for her poster at the Neuroethology meetings in Baltimore Maryland.
- Dr. Kloepper's Paper "Active echolocation beam focusing in the false killer whale (*Pseudorca crassidens*)" was chosen as paper of the month by the editors of the *Journal of Experimental Biology*.
- Dr. Kloepper also received a full NSF post-doctoral fellowship to work at Brown University with Dr. Jim Simmons on the echolocation of Bats.